

Republic of Yemen
Ministry of Higher Education & Scientific Research
Emirates International University
Faculty of Medicine and Health Sciences
Department of Clinical Pharmacy



Breast Cancer Therapy in the National Oncology Center – Yemen: Descriptive Study

A graduation research project submitted in partial fulfillment of requirements of the 5th year for obtaining bachelor degree in clinical pharmacy

Submitted by Students:

Abdulhameed Ahmed Mohammed Al-Jundubi	Abdulkafy Abdullah Abdulkarem Mohammed
Amal Ali Hadi Essa	Mohammed Muneer Humaid
Fatima Khaled Mohammed Abobakr	Naif Mohammed Al-Ameri
Ahmed Ali Ahmed Manaser Al-Khubani	Zayed Ahmed Bauam
Shoib Kamil Drhim Nagi	

Under the Supervision of:

Dr. Shawqi Hussein Nagi Al-Awdi

Assistant Professor of Pharmacology and Clinical Pharmacy
Faculty of Medicine and Health Sciences – Tamar University

Dr. Abdullah Thawabah

Oncology and Radiotherapy Consultant
General Director – National Oncology Center

2019 – 2020



Dedication

This research is dedicated to our families who gave us support, not just financially but also physically and morally. They also supported us with their love, prayers, caring and sacrifices for educating and preparing us for our future.

This research is also dedicated to all the National Oncology Center crew who helped and supported us throughout our journey of making this research. We highly appreciate the time and efforts that they have given to us especially to the respondent who spared a little of their time in sharing their experience. Without all your endeavors, we will not be able to accomplish our mission.

The researchers



Acknowledgment

First and foremost, praises and thanks to the god, the Almighty, for his showers of blessings throughout our research work to complete the research successfully.

We would like to express our deep and sincere gratitude to our research supervisor Dr. Shawqi Al-Awdi Assistant Professor of Pharmacology and Clinical Pharmacy, for giving us the opportunity to do the research and providing invaluable guidance throughout this research. His dynamism, vision, sincerity and motivation have deeply inspired us. It was a great privilege and honor to work and study under his guidance, we are extremely grateful for what he offered to us.

We would like to thank Dr. Abdullah Thawabah, general director – National Oncology Center, and his crew for their support and contribution throughout this journey.

We also would like to thank our head of department Dr. Mokhtar Al-Ghorafi for his continued support, encouragement and patience throughout the 6 years. We are extremely grateful to our professors who provided us the knowledge and shared their experience throughout the 6 years.

The researchers

Contents

Subject	Page
Contents.....	i
List of tables.....	ii
List of figures.....	iv
Abstract.....	vi
Introduction and aim of the work.....	1
Review of Literature.....	3
Subjects and Methods	26
Results.....	28
Discussion.....	55
Summary and conclusions.....	58
References.....	61
Arabic Summary	----



List of Tables

Title	Page
Table i: Breast imaging and reporting system	16
Table ii: American joint commission on cancer guidelines-tumor node metastasis classification	21
Table iii: Clinical staging-American joint commission on cancer guidelines	23
Table iv: Common combination regimens for treatment of breast cancer	24
Table 1: Age distribution for a sample of breast cancer patients	28
Table 2: Marital status distribution among a sample of breast cancer patients	29
Table 3: Ethnicity distribution among a sample of breast cancer patients	30
Table 4: Region distribution among a sample of breast cancer patients	31
Table 4: Occupation distribution among a sample of breast cancer patients	33
Table 6: Education among a sample of breast cancer patients	34
Table 7: Smoking habit among a sample of breast cancer patients	35
Table 8: Khat habituation among a sample of breast cancer patients	36
Table 9: History of hormonal contraceptive use among a sample of breast cancer patients	37
Table 10: Age at first menarche among a sample of breast cancer patients	38



Table 11: Age at first delivery among a sample of breast cancer patients	39
Table 12: Average number of children among a sample of breast cancer patients	40
Table 13: Estrogen status among a sample of breast cancer patients	41
Table 14: Progesterone status among a sample of breast cancer patients	42
Table 15: HER-2 status among a sample of breast cancer patients	43
Table 16: Invasive history among a sample of breast cancer patients	44
Table 17: TNM clinical staging among a sample of breast cancer patients	45
Table 18: Mode of treatment among a sample of breast cancer patients	46
Table 19: Type of chemotherapy among a sample of breast cancer patients	47
Table 20: Number of chemotherapy cycles among a sample of breast cancer patients	49
Table 21: Types of surgery undertaken for a sample of breast cancer patients	50
Table 22: Hormonal therapy among a sample of breast cancer patients	51
Table 23: Biological therapy among a sample of breast cancer patients	52
Table 24: Post-radiation therapy complications among a sample of breast cancer patients	53
Table 25: Post-surgical complications among a sample of breast cancer patients	54

List of Figures

Title	Page
Figure1: Age distribution for a sample of breast cancer patients	28
Figure 2: Marital status distribution among a sample of breast cancer patients	29
Figure 3: Ethnicity distribution among a sample of breast cancer patients	30
Figure 4: Region distribution among a sample of breast cancer patients	32
Figure 5: Occupation distribution among a sample of breast cancer patients	33
Figure 6: Education among a sample of breast cancer patients	34
Figure 7: Smoking habit among a sample of breast cancer patients	35
Figure 8: Khat habituation among a sample of breast cancer patients	36
Figure 9: History of hormonal contraceptive use among a sample of breast cancer patients	37
Figure 10: Age at first menarche among a sample of breast cancer patients	38
Figure 11: Age at first delivery among a sample of breast cancer patients	39
Figure 12: Average number of children among a sample of breast cancer patients	40
Figure 13: Estrogen status among a sample of breast cancer patients	41
Figure 14: Progesterone status among a sample of breast cancer patients	42



Figure 15: HER-2 status among a sample of breast cancer patients	43
Figure 16: Invasive history among a sample of breast cancer patients	44
Figure 17: TNM clinical staging among a sample of breast cancer patients	45
Figure 18: Mode of treatment among a sample of breast cancer patients	46
Figure 19: Type of chemotherapy among a sample of breast cancer patients	48
Figure 20: Number of chemotherapy cycles among a sample of breast cancer patients	49
Figure 21: Types of surgery undertaken for a sample of breast cancer patients	50
Figure 22: Hormonal therapy among a sample of breast cancer patients	51
Figure 23: Biological therapy among a sample of breast cancer patients	52
Figure 24: Post-radiation therapy complications among a sample of breast cancer patients	53
Figure 25: Post-surgical complications among a sample of breast cancer patients	54

Abstract

Breast Cancer Therapy in the National Oncology Center – Yemen: Descriptive Study

Background: The present study was aimed to assess treatment patterns among breast cancer patients at the National Oncology Cancer, Sana'a.

Methods: A hospital-based prospective cross-sectional study was conducted by collecting data from medical records of breast cancer patients who attended the breast cancer unit at the National Oncology Center in Sana'a during the period of November 1, 2019 to February 29, 2020.

Results: A total number 300 patients with breast cancer were mostly from Sana'a and Taiz governorates with a positive contraceptive history among 42.3% of patients. Out of all patients, 40.3% % were estrogen positive, 36% were progesterone positive and 51.7% were HER-2 positive. All patients have invasive form of breast cancer. Most of the patients (95%) have ductal invasive breast cancer as compared to 1.7% who have invasive lobular carcinoma. About 39% of the breast cancer patients were at stage 3, while 27.7% were at stage 4 and 25.4% were at stage 2. The most common treatment for breast cancer among the included breast cancer patients was combined surgery and chemotherapy (39%), then the standard method with surgery, radiotherapy, and chemotherapy in 32.7% of the patients. Chemotherapy alone represented 24.7% of the patients while chemotherapy along with radiotherapy represented 3.7% of the patients. Surgical procedures were undertaken for 71.4% of breast cancer patients. About 42.6% of patients with human epidermal growth factor receptor-2 (HER-2) positive breast cancer have not received biological therapy despite their positive receptor status.

Conclusions: All patients have invasive form of breast cancer which denotes that patients are usually diagnosed lately. Thus, screening of breast cancer is advised especially for patients with risk factors. Large proportion (42.6%) with human epidermal growth factor receptor-2 (HER-2) positive breast cancer patients have not received biological therapy despite their positive receptor status.



Chapter: 1

Introduction and Aim of the study



1.1 Introduction

Breast cancer is a leading cause of death worldwide, and ranks as the fifth cause of death from all cancers, and the most common cause of cancer death in women in both developing and developed countries. Prevalence of breast cancer alone accounts for 25% of all cancer cases and 15% of all cancer deaths among females (**Torre et al, 2015**).

Breast cancer is classified by whether the cancer started in the ducts or lobules, whether it grows or spread through the duct or lobule and how the cancer cells look under a microscope. It is broadly grouped in to those that are still in the breast lobules or ducts called noninvasive or carcinoma in situ and those that have started to grow and spread beyond the walls of the ducts or lobules called invasive carcinoma. Breast cancer stage is defined on the basis of the primary tumor extent and size (T1–4), presence and extent of lymph node involvement (N1–3), and presence or absence of distant metastases (M0–1) (**Dipiro et al, 2008**). Treatment of breast cancer is dependent on disease stage, histologic and molecular subtypes and menopausal status (**Di Leo et al, 2015**). Treatment of breast cancer includes surgery, radiation therapy, or both, and systemic treatment with chemotherapy, endocrine therapy, biologic therapy or combinations of these. The selection of various local or systemic therapies are based on several



prognostic and predictive factors including tumor histology, clinical and pathologic characteristics of the primary tumor, axillary lymph node status, tumor hormone receptor (estrogen receptor (ER)/ progesterone receptor (PR)) content, tumor human epidermal receptor (HER2) status, presence or absence of detectable metastatic disease, patient comorbid conditions, patient age, and menopausal status (EMRO, 2015).

1.2. Aims of the study:

1.2.1.General objective

The present study was aimed to describe treatment patterns among breast cancer patients at the National Oncology Cancer.

1.2.2. Specific objectives

- To describe the clinical characteristics of breast cancer patients.
- To describe treatment modalities of breast cancer.



Chapter: 2

Review of Literature



2. Literature Review

2.1 Introduction:

Breast cancer is the most common cancer and also the leading cause of cancer mortality in women worldwide. Approximately 1.38 million new breast cancer cases were diagnosed in 2008 with almost half of all breast cancer cases and nearly 60% of deaths occurring in lower income countries **(Ferly et al , 2010)** .There is a large variation in breast cancer survival rates around the world, with an estimated 5-year survival of 80% in high income countries to below 40% for low income countries **(Coleman et al, 2008)**.

Low and middle income countries face resource and infrastructure constraints that challenge the goal of improving breast cancer outcomes by early detection, diagnosis and treatment **(Anderson et al, 2008)** . In high income countries like the United States, approximately 232340 women will be diagnosed and 39620 will die of breast cancer in 2013 **(Seigel et al, 2013)**.

For an American woman, the lifetime risk of developing breast cancer is 12.38% or 1 in 8 **(Seigel et al, 2013)**. The significant decrease in breast cancer-related mortality in the United States from 1975 to 2000 is attributed to continued improvement in both screening mammography and treatment **(Berry et al, 2005; Ries et al, 1975-2005)**. According to the World Health



Organization, improving breast cancer outcome and survival by early detection remains the cornerstone of breast cancer control.

2.2 Risk factor prediction:

Age, reproductive factors, personal or family history of breast disease, genetic pre-disposition and environmental factors have been associated with an increased risk for the development of female breast cancer.

2.2.1 Age

The risk of developing breast cancer increases with age. By using the Surveillance, Epidemiology, and End Results (SEER) database, the probability of a woman in the United states developing breast cancer is a lifetime risk of 1 in 8; 1 in 202 from birth to age 39 years of age, 1 in 26 from 40-59 years, and 1 in 28 from 60-69 years (Siegel et al, 2013).

2.2.2 Personal history

A personal history of breast cancer is also a significant risk factor for the development of a second ipsilateral or contralateral breast cancer. In fact, the most common cancer amongst breast cancer survivors is a metachronous contralateral breast cancer (Curtis et al, 2006). Factors associated with an increased risk of a second breast cancer include an initial diagnosis of DCIS,

stage IIB, hormone receptor negative cancers, and young age (**Buist et al, 2010**).

2.2.3 Breast pathology

Proliferative breast disease is associated with an increased risk of breast cancer. Proliferative breast lesions without atypia, including usual ductal hyperplasia, intraductal papillomas, sclerosing adenosis and fibroadenomas confer only a small increased risk of breast cancer development, approximately 1.5-2 times that of the general population (**Hartmann et al, 2005**). Atypical hyperplasia including both ductal and lobular, usually incidentally found on screening mammography, confers a substantial increased risk of breast cancer. Women with atypia have an approximately 4.3 times greater risk of developing cancer compared to the general population (**Hartmann et al, 2005; Doupont et al, 1993**).

2.2.4 Family history

A woman's risk of breast cancer is increased if she has a family history of the disease. In the Nurses' Health Study follow-up, women with a mother diagnosed before age 50 had an adjusted relative risk of 1.69 and women with a mother diagnosed at 50 or older had a relative risk of 1.37 compared to women without a family history of breast cancer. A history of a

sister with breast cancer also demonstrated an increased relative risk of 1.66 if the diagnosis was made prior to age 50 and a relative risk of 1.52 if diagnosed after age 50 compared to patients without a family history (**Colditz et al, 2012**). The highest risk is associated with increasing number of first degree relatives diagnosed with breast cancer at a young age (under age 50). Compared with women who had no affected relative, women who had one, two or three or more affected first degree relatives had risk ratios of 1.80, 2.93 and 3.90, respectively (**Collaborative Group on Hormonal Factors in Breast , 2001**).

2.2.5 Genetic predisposition

Approximately 20%-25% of breast cancer patients have a positive family history but only 5%-10% of breast cancer cases demonstrate an autosomal dominant inheritance (**Lynch and Lynch, 1986; Margolin et al, 2006**). Genetic predisposition alleles have been described in terms of clinical significance (**Lalloo and Evans, 2012**). High-risk predisposition alleles conferring a 40%-85% lifetime risk of developing breast cancer include BRCA1 and BRCA2 mutations, mutations in TP53 gene resulting in Li-Fraumeni syndrome, PTEN resulting in Cowden syndrome, STK11 causing Peutz-Jegher's syndrome, Neurofibromatosis (NF1) and (CDH-1) E-Cadherin (**Sharif et al, 2007**). Half of the breast cancer predisposition

syndromes are associated with mutations in BRCA1 and BRCA2. Women with BRCA1 or BRCA2 deleterious mutations have a significantly higher risk of developing breast cancer. Lifetime breast cancer risk ranges from 65% to 81% for BRCA1 mutation carriers and 45% to 85% for BRCA2 carriers (Ford et al, 1998). Moderate risk genes including homozygous ataxia-telangiectasia (ATM) mutations (Thompson et al, 2005), somatic mutations in tumor suppressor gene CHEK2, and BRCA1 and BRCA2 modifier genes BRIP1 (Seal et al, 2006) and PALB2 (Wong et al, 2011) confer a 20%-40% lifetime risk of breast cancer. Numerous low risk common alleles have been identified largely through genome-wide association studies (Lalloo and Evans , 2012) and the clinical application in the presence of these mutations is yet to be determined.

2.4 Endogenous hormone exposure and reproductive factors:

The cycles of endogenous estrogen levels throughout a woman's lifetime have implications for the development of or the protection against breast cancer.

2.4.1 Early menarche

Early age at menarche is a risk factor among both pre and postmenopausal women for developing breast cancer. Delay in menarche by two years is associated with corresponding risk reduction of 10% (Hsieh et al,

1990). Within the European Prospective Investigation into Cancer and Nutrition cohort, women who had early menarche (≤ 13 years) demonstrated a nearly twofold increase in risk of hormone receptor positive tumors (Ritte et al, 2012).

2.4.2 Parity and age at first full term pregnancy

Nulliparous women are at an increased risk for the development of breast cancer compared to parous women. Young age at first birth has an overall protective effect, whereas relatively advanced age at first birth confers a relative risk of breast cancer greater than that of a nulliparous woman. Compared to nulliparous women the cumulative incidence of breast cancer in women experiencing their first birth at age 20, 25, and 35 years was 20% lower, 10% lower and 5% higher, respectively (Rosner et al, 1994).

2.4.3 Breast feeding

Evidence suggests that breast feeding has a protective effect against the development of breast cancer. Breast feeding may delay return of regular ovulatory cycles and decrease endogenous sex hormone levels. It has been estimated that there is a 4.3% reduction for every one-year of breast feeding (Collaborative Group on Hormonal Factors , 2002).

2.4.4 Testosterone

High endogenous sex hormone levels increase the risk of breast cancer in both premenopausal and postmenopausal women. High levels of circulating testosterone in postmenopausal women have been linked to increased risk of developing breast cancer [relative risk (RR), 2.86-3.28] (Sieri et al, 2009).

2.4.5 Age at menopause


Later onset of menopause has also been associated with increased breast cancer risk. Every year delay in the onset of menopause confers a 3% increase in risk and every five year delay in the onset of menopause confers a 17% increase in risk of breast cancer (Hsieh et al, 1990; Kelsey et al, 1993).

2.5 Exogenous hormone exposure:

Evidence suggests a relationship between the use of hormone replacement therapy (HRT) and breast cancer risk. Breast cancers related to HRT use are usually hormone receptor positive. When compared with patients who do not use HRT, breast cancer risk is higher in HRT users (Lahmann et al, 2004). An international meta-analysis examining the risk of breast cancer with HRT found that in women who did not use HRT, RR

increased by a factor of 1.028 for each year older at menopause, comparable to the relative risk of 1.023 per year in women who use HRT or for those who ceased to use HRT up to four years previously (**Lancet, 1997**).

In the Woman's Health Initiative randomized control trial, combined estrogen plus progestin in postmenopausal women with an intact uterus significantly increased the risk of breast cancer, delayed breast cancer detection and diagnosis, and significantly increased breast cancer mortality. The study was terminated early because of increased mortality in the combined estrogen plus progestin group. By contrast, the use of estrogen alone by postmenopausal women without a uterus did not interfere with breast cancer detection and statistically significantly decreased the risk of breast cancer (**Anderson et al, 2003**). Data from the Nurses' Health Study, however, suggest that women who use unopposed postmenopausal estrogen increase their risk of breast cancer by 23% at age 70 (**Colditz and Rosner , 2000**). Timing and duration of HRT seem to be important factors associated with breast cancer risk as well. Breast cancer risk from exogenous hormone exposure is inversely associated with time from menopause. Women initiating hormone therapy closer to menopause have a higher breast cancer risk (**Chlebowski et al, 2013**). Long term (> 5 years) combined HRT use has been associated with the highest risk whereas short-term use of



combined estrogen-progestin therapy does not appear to confer a significantly increased risk (RR = 1.023 per year) (**lancet, 1997**).

2.6 Lifestyle factors

Modifiable risk factors including the excessive use of alcohol, obesity and physical inactivity account for 21% of all breast cancer deaths worldwide (**Danaei et al, 2005**).

2.6.1 Alcohol consumption

Alcohol consumption has been associated with increased breast cancer risk that is statistically significant at levels as low as 5.0 to 9.9 g per day, equivalent to 3 to 6 drinks per week. Binge drinking, but not frequency of drinking, was associated with breast cancer risk after controlling for cumulative alcohol intake. Alcohol intake both earlier and later in adult life was independently associated with risk (**Chen et al, 2011**).

2.6.2 Physical activity

Consistent physical activity has been shown to reduce the risk of breast cancer in a dose dependent manner, with modest activity conferring a 2% decrease in risk and vigorous activity a 5% decrease in risk (**Wu et al, 2013**).



2.6.3 Obesity

Obesity, specifically in postmenopausal women, has also been shown to increase a woman's risk of breast cancer. In the EPIC multicenter prospective cohort study, postmenopausal women who did not use HRT had elevated breast cancer risk with increasing weight, body mass index (BMI) and hip circumference **(Lahmann et al, 2004)**. In this cohort, multivariate relative risk was 1.28 for overweight women (BMI 25.0-29.9) and obese women (BMI > 30.0) compared to women in the normal weight range. Lean women on HRT are incongruously at an increased risk of breast cancer (RR = 2.04) compared to their overweight (1.93) and obese (1.39) counterparts **(Lahmann et al, 2004)**. Insulin resistance and hyperinsulinemia have been studied as a risk factor for the comorbidities associated with obesity including cardiovascular disease and diabetes. Insulin has anabolic effects on cellular metabolism and insulin receptor overexpression has been demonstrated in human cancer cells **(Milazzo et al, 1992)**. Hyperinsulinemia has been shown to be an independent risk factor for breast cancer in nondiabetic postmenopausal women and may help to explain the relationship between obesity and breast cancer **(Gunter et al, 2009)**.



2.6.4 Radiation

Radiation exposure from various sources including medical treatment and nuclear explosion increases the risk of breast cancer. Radiation to the chest wall for treatment of childhood cancer increases the risk of breast cancer linearly with chest radiation dose (**Henderson et al, 2010**). Survivors of childhood cancers who received therapeutic radiation are at a dose dependent risk for the development of breast cancer, and those treated for Hodgkin's disease are at highest risk ($RR = 7$) (**Guibout et al, 2005**). Radiation effects on the development of female breast cancer have also been demonstrated in Japan post nuclear attack on Hiroshima and Nagasaki (**Preston et al, 2007**) and positively correlate with age younger than 35 years at time of exposure. The incidence of breast cancer has also increased in areas of Belarus and Ukraine. A significant two fold increase was seen in the most contaminated areas around Chernobyl following the nuclear accident and manifest in women who were younger at the time of the exposure (**Pukkala et al, 2006**).

2.7 Screening

2.7.1 Breast self- and clinical breast examination

Utility of the breast self-examination (BSE) is controversial as the benefit in terms of decreased mortality has not been demonstrated (**Kösters and Göttsche, 2003**). Most clinicians encourage women to perform monthly BSE to become familiar with their normal anatomy and empower them with regards to their own healthcare (**McCready et al, 2005**). The 2013 NCCN guidelines recommend annual clinical breast examination (CBE) for women of average risk > 40 years of age as well as BSE to develop and exhibit breast self-awareness (**National Comprehensive Cancer Network, 2013**).

2.7.2 Mammography

One of the most important advances in the treatment of breast cancer is early detection of non-palpable masses. In the 1960's, the first randomized control trials comparing periodic mammography screening vs clinical examination demonstrated a decreased mortality by approximately one third in the experimental group. However there is still controversy regarding mortality from breast cancer in the subset of women aged 40-49 years (**Shapiro et al, 1971; Shapiro et al, 1985**). Contemporary randomized control trials have demonstrated the benefits from screening mammography

in women aged 40 to 70 years (**Freedman et al, 2004; Nyström et al, 2002**). A 2013 Cochrane Review suggests that mortality is an outcome biased toward screening, routine mammography leads to undue stress and uncertainty in the face of false-positive results with increase in total numbers of lumpectomies and mastectomies but no decrease in mortality (**Gøtzsche and Jørgensen , 2013**). Controversy surrounding mammography is related to the inherent lead time and length time biases in screening for disease. Lead time bias is an overestimation of survival among screen detected cases compared to clinically detected cases when true survival time actually remains unchanged. Length bias is an overestimation of survival time among screening-detected cases, which is caused by those slowly progressing cases that may never be clinically relevant. The 2013 NCCN guidelines recommends annual screening mammography in women ≥ 40 years of average risk and annual mammography at age 25 or individualized based on onset of cancer in proband in patients who are high risk by prediction models, known history or genetic predisposition syndrome as well as the counseling and education of risks and benefits related to participating in cancer screening (**National Comprehensive Cancer Network, 2013**).

Mammography remains the mainstay in breast cancer detection (**Smetherman , 2013**). Diagnostic mammograms are performed in women



who have a palpable mass or other symptom of breast disease, a history of breast cancer within the preceding 5 years, or have been recalled for additional imaging from an abnormal screening mammogram. Diagnostic mammograms include special views such as focal compression of one area of the breast tissue or magnification images. The breast imaging reporting and database system (BI-RADS) is the standardized method for reporting of mammographic findings (American College of Radiology, 2003). Carcinomas present as masses, asymmetries, and calcifications (Table 1).

Table 1: breast imaging and reporting system

Category	Assessment	Follow-up
0	Need additional imaging evaluation	Additional imaging needed before a category can be assigned
1	Negative	Continue annual screening mammograms (women older than 40 yr)
2	Benign	Continue annual screening mammograms (women older than 40 yr)
3	Probably benign	Initial short term follow-up (usually six month mammogram (< 2% chance of malignancy)
4	Suspicious abnormality	Biopsy should be considered (2%-95% chance of malignancy)
5	Highly suggestive of malignancy	Requires biopsy (> 95% chance of malignancy)
6	Known cancer	Biopsy-proven malignancy



2.7.3 Magnetic resonance imaging

Mammography remains the gold standard for breast imaging but magnetic resonance imaging (MRI) has become an important modality in the detection, assessment, staging, and management of breast cancer in selected patients. Screening MRI is more sensitive but less specific for the detection of cancer in high risk women. The sensitivity of MRI is 0.77-0.79 compared to mammographic sensitivity of 0.33-0.39. Specificity of MRI is 0.86-0.89 compared to mammographic specificity of 0.95 (**Warner et al, 2008; Kriege et al, 2004**). In a systematic review, MRI and mammography demonstrated a combined sensitivity and specificity of 0.94 and 0.77, respectively (**Warner et al, 2008**). The 2013 NCCN guidelines recommend patients who have increased ($> 20\%$) lifetime risk of developing breast cancer undergo annual mammography and MRI starting at age 25 or an age tailored to the risk of the patient on an individual basis. MRI is valuable in the screening of select high risk patients, patients in whom breast augmentation prevents effective screening mammography, or in patients with equivocal findings on other imaging modalities.

2.7.4 Ultrasound

There are several studies supporting the use of adjunctive screening ultrasound in high risk patients with dense breast tissue, which imparts a substantial but accepted number of false positives (**Berg et al, 2008**). No randomized controlled trials have been conducted to evaluate the impact of screening ultrasonography on breast cancer mortality rates. Whole breast ultrasound may allow the clinician to screen for breast cancers not detected by traditional mammography, especially in dense breasts where mammographic sensitivity is lower (**Kelly et al, 2010**). Single center studies have shown that the incremental detection of breast cancer by ultrasound following screening mammogram offers only marginal added benefit in women of average risk (**Gartlehner et al, 2013**).

2.10 Prognostic indicators

2.10.1 Estrogen receptor and progesterone receptor status

Estrogen receptor (ER) and progesterone receptor (PR) represent weak prognostic factors for patients with breast cancer, but these receptors are the strongest predictive factors for response to endocrine therapy. ER and PR assays should be performed on all invasive breast cancers (**Fitzgibbons et al, 2000**). Both ER and PR are assessed by

immunohistochemistry (IHC) on paraffin sections. IHC allows assessment of the expression specifically in either invasive or *in situ* cancer. Positive interpretation requires at least 1% of tumor cells showing positive nuclear staining of any intensity. Receptor negative is reported if less than 1% of tumor cells show staining of any intensity (Hammond et al, 2010). The cutoff to define positivity is 1% because patients with even 1% ER/PRpositive tumors may benefit from hormonal therapy. About 70% of all breast cancers are ER-positive and 60% to 65% of all breast cancers are PR-positive. For the patients with a “weak positive” result an Allred score helps differentiate positive from negative receptor status. The Allred score categorizes the percentage of cells (scored from 0 to 5) with the intensity (scored from 0 to 3) and adds these two scores to give a numerical score from 0 to 8 (Allred et al, 1998). A score of 0-2 was regarded as negative and 3-8 as positive.

2.10.2 HER2 protein expression and gene amplification

HER-2/neu is a proto-oncogene that encodes for a transmembrane tyrosine kinase growth receptor, and it is involved in several regulatory pathways in breast, involving proliferation, survival, cell motility, and invasion. HER2 is usually assessed by IHC. Fluorescence *in situ* hybridization (FISH) assay of HER2 expression is usually performed when



the evaluation by IHC is equivocal. HER2 is a prognostic factor for outcome in both node-negative and node-positive patients and is a predictive factor for response to certain therapies that target the HER-2/neu receptor such as trastuzumab (Herceptin), a monoclonal antibody targeted to the HER2 protein, and other newer anti-HER2 agents. Overexpression/amplification is reported in 10% to 34% of invasive breast cancers. Gene over expression and amplification and surface membrane protein expression are concordant in more than 90% of cases (Wolff et al, 2007; Hicks & Kulkarni , 2008).

2.10.3 Commercially available gene assays

OncotypeDX (Genomic Health, Inc, Redwood City, California) is a reverse transcription polymerase chain reaction-based assay that can be performed on paraffin sections. It is based on analysis of the expression of 21 genes and provides a “recurrence score” that correlates with outcome. Although it was initially used to assess prognosis in ER-positive, node-negative patients (Paik et al, 2004), data have indicated that it is an equally valuable prognostic indicator in ER-positive, node-positive patients. Another molecular profiling product is the Amsterdam 70-gene profile, Mammaprint (Agendia, Amsterdam, Netherlands), in which a microarray analysis of gene expression is used on breast cancer tissue. It is used to determine the prognosis of patients with breast cancer and can be used for all tumors,

including node-positive, *HER-2 neu*-positive, and ER/PR-negative disease (van de Vijver et al, 2002).

2.11 Staging:

After a breast cancer has been diagnosed, the patient is clinically staged using the American Joint Commission on Cancer (AJCC) guidelines (Tables 2 and 3).

Table 2: American Joint Commission on Cancer guidelines–tumor node metastasis classification:

Primary tumor (T)	
TX	Primary tumor cannot be assessed
T0	No evidence of primary tumor
Tis	Carcinoma in situ
Tis (DCIS)	Ductal carcinoma in situ
Tis (LCIS)	Requires biopsy (> 95% chance of malignancy)
Tis (Paget's)	Paget's disease of the nipple
T1	Tumor \leq 20 mm in greatest dimension
T1mi	Tumor \leq 1 mm in greatest dimension
T1a	Tumor > 1 mm but \leq 5 mm in greatest dimension
T1b	Tumor > 5 mm but \leq 10 mm in greatest dimension
T1c	Tumor > 10 mm but \leq 20 mm in greatest dimension
T2	Tumor > 20 mm but \leq 50 mm in greatest dimension
T3	Tumor > 50 mm in greatest dimension
T4	Tumor of any size with direct extension to the chest wall and/or to the skin (ulceration or skin nodules)
T4a	Extension to the chest wall, not including only pectoralis muscle adherence/invasion

T4b	Ulceration and/or ipsilateral satellite nodules and/or edema (including peau d'orange) of the skin, which do not meet the criteria for inflammatory carcinoma
T4c	Both T4a and T4b
T4d	Inflammatory carcinoma
Regional lymph nodes (N)	
NX	Regional lymph nodes cannot be assessed (for example, previously removed)
N0	No regional lymph node metastases
N1	Metastases to movable ipsilateral level I , II axillary lymph node(s)
N2	Metastases in ipsilateral level I , II axillary lymph nodes that are clinically fixed or matted; or in clinically detected ipsilateral internal mammary nodes in the absence of clinically evident axillary lymph node metastases Metastases in ipsilateral level I , II axillary lymph nodes fixed to one another (matted) or to other structures
N2a	Metastases only in clinically detected ipsilateral internal mammary nodes and in the absence of clinically evident level I, II axillary lymph node metastases
N2b	
N3	Metastases in ipsilateralinfraclavicular (level III axillary) lymph node(s) with or without level I , II axillary lymph node involvement; or in clinically detected ipsilateral internal mammary lymph node(s) with clinically evident level I , II axillary lymph node metastases; or metastases in ipsilateral supraclavicular lymph node(s) with or without axillary or internal mammary lymph node involvement
N3a	Metastases in ipsilateralinfraclavicular lymph node(s)
N3b	Metastases in ipsilateral internal mammary lymph node(s) and axillary lymph node(s)

N3c	Metastases in ipsilateral supraclavicular lymph node(s)
Distant metastasis (M)	
M0	No clinical or radiographic evidence of distant metastases
cM0(i+)	No clinical or radiographic evidence of distant metastases, but deposits of molecularly or microscopically detected tumor cells in circulating blood, bone marrow, or other nonregional nodal tissue that are no larger than 0.2 mm in a patient without symptoms or signs of metastases
M1	Distant detectable metastases as determined by classic clinical and radiographic means and/or histologically proven larger than 0.2 mm

Table 3: Clinical Staging-American Joint Commission on Cancer Guidelines

Stage 0	Tis	N0	M0
Stage I A	T1	N0	M0
Stage I B	T0	N1mi	M0
	T1	N1mi	M0
Stage II A	T0	N1	M0
	T1	N1	M0
	T2	N0	Mo
Stage II B	T2	N1	M0
	T3	N0	M0
Stage III A	T0	N2	M0
	T1	N2	M0
	T2	N2	M0
	T3	N1	M0
	T3	N2	M0
Stage III B	T4	N0	M0
	T4	N1	M0
	T4	N2	M0
Stage III C	Any T	N3	M0
Stage IV	Any T	Any N	M1

2.12 Management:

Table 3: Common Combination Regimens for Treatment of Breast Cancer

Regimen	Dose and Schedule	Frequency	Cycles
TAC			
T - Docetaxel	75 mg/m ² IV day 1	Every 21 days	6
A – Doxorubicin	50 mg/m ² IV day 1		
C - Cyclophosphamide	500 mg/m ² IV day 1		
AC ⇒Taxol (T) (conventional regimen)			
Doxorubicin	60 mg/m ² IV day 1	Every 21 days	4
Cyclophosphamide	600 mg/m ² IV day 1		
<i>Followed by</i>			
Paclitaxel	175 mg/m ² IV day 1	Every 21 days	4
Dose-dense			
Doxorubicin	60 mg/m ² IV day 1	Every 14 days	4
Cyclophosphamide	600 mg/m ² IV day 1		
<i>Followed by</i>			
Paclitaxel	175 mg/m ² IV day 1	Every 14 days	4
Metronomic regimen			
Doxorubicin	20 mg/m ² IV day 1	Every week	12
Cyclophosphamide	50 mg/m ² PO	Every day	
<i>Followed by</i>			
Paclitaxel	80 mg/m ² IV day 1	Every week	12
AC ⇒T + H (trastuzumab [Herceptin])			
Trastuzumab dosage: 4 mg/kg IV load, then 2 mg/kg weekly with paclitaxel, then give 6 mg/kg IV every 3 weeks for 40 weeks			
NOTE: Trastuzumab to be added to a weekly paclitaxel regimen in HER2-positive breast cancer patients			
FEC100			
5-Fluorouracil (5-FU)	500 mg/m ² IV day 1	Every 21 days	6
Epirubicin	100 mg/m ² IV day 1		
Cyclophosphamide	500 mg/m ² IV day 1		
FAC			
5-FU	600 mg/m ² IV day 1	Every 21 days	4
Doxorubicin	60 mg/m ² IV day 1		

Cyclophosphamide	600 mg/m ² IV day 1		
5-FU	500 mg/m ² IV days 1 and 8	Every 28 days	6
Doxorubicin	30 mg/m ² IV days 1 and 8		
Cyclophosphamide	100 mg/m ² PO days 1-14		
CMF (Bonadonna regimen)			
Cyclophosphamide	100 mg/m ² PO days 1-14	Every 28 days	6
Methotrexate	40 mg/m ² IV days 1 and 8		
5-FU	600 mg/m ² IV days 1 and 8		
Metronomic regimen			
Cyclophosphamide	50 mg/m ² PO days 1-7	Weekly	24
Methotrexate	15 mg/m ² IV		
5-FU	300 mg/m ² IV		
TC			
Taxotere (Docetaxel)	75 mg/m ² IV day 1	Every 21 days	4
Cyclophosphamide	600 mg/m ² IV day 1		
TCH			
Docetaxel	75 mg/m ² IV day 1	Every 21 days	6
Carboplatin	AUC* 6 IV day 1		
Trastuzumab	8 mg/kg loading dose IV followed by 6 mg/kg/wk q3wk		17
TCH-P			
Docetaxel	75 mg/m ² IV day 1	Every 21 days	6
Carboplatin	AUC*6 IV day 1		
Trastuzumab	8 mg/kg loading dose IV followed by 6 mg/kg		17
Pertuzumab	840 mg loading dose IV followed by 420 mg on subsequent doses		6

*AUC = systemic exposure.

<https://emedicine.medscape.com/article/1946040-overview#a3>

Chapter: 3

Subjects and Methods



3. Subjects and Methods:

3.1 Study protocol:

The study was a hospital-based prospective cross-sectional study by collecting data who were female older than 18 years who visited the National Oncology Center from between November 1, 2019 and February 29, 2020.

3.2 Study population:

The source population constitutes all medical records of breast cancer patients who attended the breast cancer unit at the National Oncology Center, Sana'a. The study population constitutes all medical records of breast cancer patients who attended the center that fulfill the inclusion criteria of the study.

3.3. Inclusion and exclusion criteria

Inclusion

- Medical charts of female breast cancer patients
- Age ≥ 18 years.
- Patient attended to the national oncology center between November 1, 2019 and February 29, 2020.

Exclusion

- Patients who were diagnosed with other cancer.



3.4. Sample size

All breast cancer patients who attended the breast cancer unit of the National Oncology Center from November 1, 2019 to February 29, 2020 and fulfilled the inclusion criteria of the study were included in the study. As a result, there were 300 breast cancer patients who newly visited the oncology center and fulfilled the inclusion criteria and hence were included in the study.

3.5 Statistical analysis:

Statistical analysis was performed using Statistical Package for Social Science software (SPSS, version 15) and Microsoft office Excel 2010 was used for data processing and statistical analysis. The chi-squared test was used for the assessment of association between the variables studied. The p-value of less than 0.05 was significant statistically.

Chapter: 4

Results



4. Results

4.1. Demographic data:

4.1.1 Age distribution of the patients:

Total number of sample enrolled in the present study was 300 patients with breast cancer. All patients were females. Patients' age ranged between 21 and 80 years. The present study observed that most cases of breast cancer patients aged between 30-59 years (78.4%). The age distribution of the sample is represented in **table (1)** and graphically illustrated in **Figure (1)**.

Table (1): Age distribution for a sample of breast cancer patients:

	Frequency	Percent
20-29 years	22	7.3
30-39 years	89	29.7
40-49 years	86	28.7
50-59 years	60	20.0
60-69 years	31	10.3
70-80 years	12	4.0
Total	300	100.0




Figure (1): Age distribution for a sample of breast cancer patients

4.1.2 Marital status of the patients:

The majority of enrolled patients (73.3%) were married. Widowed patients were about 13.3% while divorced patients were 4.7%. In the other hand, single patients were about 8.7%. The distribution of the sample is represented in **table (2)** and graphically illustrated in **Figure (2)**.

Table (2): Marital status distribution among a sample of breast cancer patients:

	Frequency	Percent
married	220	73.3
single	26	8.7
divorced / separated	14	4.7
widowed	40	13.3
Total	300	100.0




Figure (2): Marital status distribution among a sample of breast cancer patients

4.1.3 Ethnicity of the patients:

Almost all breast cancer patients (98%) were of Caucasian ethnicity while 2% were African race. The distribution of the sample is represented in **table (3)** and graphically illustrated in **Figure (3)**.

Table (3): Ethnicity distribution among a sample of breast cancer patients:

	Frequency	Percent
Caucasian	294	98.0
African	6	2.0
Total	300	100.0




Figure (3): Ethnicity distribution among a sample of breast cancer patients

4.1.4 Residence of the patients:

Most patients of breast cancer (31.3%) were from Sana'a and from Taiz (15.7%) governorates. The second common places of breast cancer patients were Ibb (10.3%), Thamar (8.7%), and Hodeidah (6.7%) governorates. Other governorates ranked lower. The distribution of the sample is represented in **table (4)** and graphically illustrated in **Figure (4)**.

Table (4): Region distribution among a sample of breast cancer patients:

	Frequency	Percent
Sana'a	94	31.3
Taiz	47	15.7
Thamar	31	10.3
Ibb	26	8.7
Hodeidah	20	6.7
Saada	12	4.0
Amran	11	3.7
Albaidha	10	3.3
Hajah	9	3.0
Almahweet	9	3.0
Rimah	7	2.3
Aldhalae	7	2.3
Aden	7	2.3
Lahj	3	1.0
Aljouf	2	.7
Socotra	2	.7
Hadhrmoot	1	.3
Mareb	1	.3
Mahrah	1	.3
Total	300	100.0




Figure (4): Region distribution among a sample of breast cancer patients

4.1.5 Occupation of the patients:

Most breast cancer patients (82.3%) were housewives. Peasant women were about 10.7% while employed patients represented 4.7%. The distribution of the sample is represented in **table (5)** and graphically illustrated in **Figure (5)**.

Table (4): Occupation distribution among a sample of breast cancer patients:

	Frequency	Percent
peasant	32	10.7
private	7	2.3
employed	14	4.7
house wife	247	82.3
Total	300	100.0




Figure (5): Occupation distribution among a sample of breast cancer patients

4.1.6 Education of the patients:

Most breast cancer patients (61.7%) were illiterate. Patients who have primary school education were 18.7%, and patients with secondary school certificate were 10.7%, while patients who had a university degree were 8%. The distribution of the sample is represented in **table (6)** and graphically illustrated in **Figure (6)**.

Table (6): Education among a sample of breast cancer patients:

	Frequency	Percent
no	185	61.7
primary school	56	18.7
secondary school	32	10.7
university	24	8.0
other	3	1.0
Total	300	100.0




Figure (6): Education among a sample of breast cancer patients

4.1.7 Smoking habit distribution among patients:

Smoking habit was found less popular among breast cancer patients in about 18.7% while non-smokers were 81.3%. The distribution of the sample is represented in **table (7)** and graphically illustrated in **Figure (7)**.

Table (7): Smoking habit among a sample of breast cancer patients:

	Frequency	Percent
No smoking	244	81.3
Yes smoking	56	18.7
Total	300	100.0




Figure (7): Smoking habit among a sample of breast cancer patients

4.1.8 Khat habituation among patients:

Khat habituation was found among 53.7% breast cancer patients while non-khat chewers were 46.3%. The distribution of the sample is represented in **table (8)** and graphically illustrated in **Figure (8)**.

Table (8): Khat habituation among a sample of breast cancer patients:

	Frequency	Percent
No khat chewing	139	46.3
Yes khat chewing	161	53.7
Total	300	100.0




Figure (8): Khat habituation among a sample of breast cancer patients

4.2 Reproductive risk factors:

4.2.1 Hormonal contraceptive history among patients:

History of contraceptive use was found among 42.3% of breast cancer patients while 57.7% of the patients stated there had not used any contraceptives during their life. The distribution of the sample is represented in **table (9)** and graphically illustrated in **Figure (9)**.

Table (9): History of hormonal contraceptive use among a sample of breast cancer patients:

	Frequency	Percent
no contraceptive history	173	57.7
yes contraceptive history	127	42.3
Total	300	100.0




Figure (9): History of hormonal contraceptive use among a sample of breast cancer patients

4.2.2 Age at first menarche:

Age at first menarche was less than 11 years in about 10.3% of breast of breast cancer patients while 89.7% of the patients aged at first menarche between 12 and 18 years. The distribution of the sample is represented in **table (10)** and graphically illustrated in **Figure (10)**.

Table (10): Age at first menarche among a sample of breast cancer patients:

	Frequency	Percent
9-11 years	31	10.3
12-18 years	269	89.7
Total	300	100.0




Figure (10): Age at first menarche among a sample of breast cancer patients

4.2.3 Age at first delivery:

About 18% of breast cancer patients were nulliparous as the either had not children despite marriage or because they are single. Age at first delivery ranged between 12 and 42 years. Most patients 71.3% had got their children while they aged 12-29 years old. About 6% of the patients had their children between 30-34 years while 4.7% of the patients had their children after age 35 years old. The distribution of the sample is represented in **table (11)** and graphically illustrated in **Figure (11)**.

Table (11): Age at first delivery among a sample of breast cancer patients:

	Frequency	Percent
no children	54	18.0
12-19 years	90	30.0
20-29 years	124	41.3
30-34 years	18	6.0
35-42 years	14	4.7
Total	300	100.0




Figure (11): Age at first delivery among a sample of breast cancer patients

4.2.4 Number of children:

The number of children ranged between 1 and 14 for breast cancer women. Relatively large percentage (38.7%) of mothers had 3-6 children followed by mothers having 7-10 children (18.3%) or 1-2 children (16%). In the other hand, 9% of enrolled mothers had 11-14 children. The distribution of the sample is represented in **table (12)** and graphically illustrated in **Figure (12)**.

Table (12): Average number of children among a sample of breast cancer patients:

	Frequency	Percent
no children	54	18.0
1-2 children	48	16.0
3-6 children	116	38.7
7-10 children	55	18.3
11-14 children	27	9.0
Total	300	100.0




Figure (12): Average number of children among a sample of breast cancer patients

4.3 Clinical characteristics of the study patients:

4.3.1 Estrogen receptor status:

Out of a total 300 breast cancer patients, 40.3% were estrogen positive, and 59.7% were estrogen negative. The distribution of the sample is represented in **table (13)** and graphically illustrated in **Figure (13)**.

Table (13): Estrogen status among a sample of breast cancer patients:

	Frequency	Percent
negative	179	59.7
positive	121	40.3
Total	300	100.0




Figure (13): Estrogen status among a sample of breast cancer patients

4.3.2 Progesterone receptor status:

Out of a total 300 breast cancer patients, 36% were progesterone positive, and 64% were progesterone negative. The distribution of the sample is represented in **table (14)** and graphically illustrated in **Figure (14)**.

Table (14): Progesterone status among a sample of breast cancer patients:

	Frequency	Percent
negative	192	64.0
positive	108	36.0
Total	300	100.0




Figure (14): Progesterone status among a sample of breast cancer patients

4.3.3 Human epidermal growth factor receptor-2 (HER 2) status:

Out of a total 300 breast cancer patients, 51.7% were HER-2 positive, and 48.3% were HER-2 negative. The distribution of the sample is represented in **table (15)** and graphically illustrated in **Figure (15)**.

Table (15): HER-2 status among a sample of breast cancer patients:

	Frequency	Percent
negative	145	48.3
positive	155	51.7
Total	300	100.0




Figure (15): HER-2 status among a sample of breast cancer patients

4.3.4 Invasive history:

About 95% of the patients have invasive ductal form of breast cancer as compared to 1.7% who have invasive lobular carcinoma in situ. Some patients (2%) have both ductal and lobular invasive breast cancer. The distribution of the sample is represented in **table (16)** and graphically illustrated in **Figure (16)**.

Table (16): Invasive history among a sample of breast cancer patients:

	Frequency	Percent
ductal invasive	285	95.0
lobular invasive	5	1.7
ductal and lobular invasive	6	2.0
ductal and medullar invasive	2	.7
ductal and tubular invasive	2	.7
Total	300	100.0

Invasive histology




Figure (16): Invasive history among a sample of breast cancer patients

4.3.5 TNM clinical staging of breast cancer:

About 39% of the breast cancer patients were at stage 3, while 27.7% were at stage 4 and 25.4% were at stage 2. In the other hand, 6.7% of breast cancer patients have not been stated their clinical stage. Only, 1.3% of the patients were at stage 1. The distribution of the sample is represented in **table (17)** and graphically illustrated in **Figure (17)**.

Table (17): TNM clinical staging among a sample of breast cancer patients:

	Frequency	Percent
stage 1	4	1.3
stage 2a	20	6.7
stage 2b	56	18.7
stage 3a	54	18.0
stage 3b	34	11.3
stage 3c	29	9.7
stage 4	83	27.7
not-stated	20	6.7
Total	300	100.0

TNM clinical staging




Figure (17): TNM clinical staging among a sample of breast cancer patients

4.5 Treatment of breast cancer:

4.5.1 Mode of treatment:

The most common treatment for breast cancer among the included breast cancer patients was combined surgery and chemotherapy (39%), then the standard method with surgery, radiotherapy, and chemotherapy in 32.7% of the patients. Chemotherapy alone represented 24.7% of the patients while chemotherapy along with radiotherapy represented 3.7% of the patients. The distribution of the sample is represented in **table (18)** and graphically illustrated in **Figure (18)**.

Table (18): Mode of treatment among a sample of breast cancer patients:

	Frequency	Percent
chemotherapy	74	24.7
radiotherapy and chemotherapy	11	3.7
chemotherapy, radiotherapy, surgery	98	32.7
surgery and chemotherapy	117	39.0
Total	300	100.0

Mode of treatment




Figure (18): Mode of treatment among a sample of breast cancer patients

4.4.2 Type of chemotherapy:

The most common combined regimens used for treatment of breast cancer were AC→T “doxorubicin + cyclophosphamide → taxol” (34.3%) followed by AC “doxorubicin + cyclophosphamide” (16.3%). FAC regimen “5-fluorouracil + doxorubicin + cyclophosphamide” was prescribed for 14%, and FAC→T regiment “5-fluorouracil + doxorubicin + cyclophosphamide → taxol” was prescribed for 13% of the patients. Other patients were treated with various or sequential chemotherapy combinations. The distribution of the sample is represented in **table (19)** and graphically illustrated in **Figure (19)**.

Table (19): Type of chemotherapy among a sample of breast cancer patients:

	Frequency	Percent
ACT	103	34.3
AC	49	16.3
FAC	42	14.0
FACT	39	13.0
ACTP or ACTGP	19	6.3
FACTP or FACTGP	19	6.3
other	29	9.7
Total	300	100.0




Figure (19): Type of chemotherapy among a sample of breast cancer patients

4.4.3 Number of chemotherapy cycles:

Chemotherapy cycles ranged from 4-18 cycles. Most patients (32%) have received 8 cycles of chemotherapy regimens. Large percent of patients received either 6 (17.7%) or 4 cycles (15.7%). The distribution of the sample is represented in **table (20)** and graphically illustrated in **Figure (20)**.

Table (20): Number of chemotherapy cycles among a sample of breast cancer patients:

	Frequency	Percent
4.00	47	15.7
6.00	53	17.7
8.00	96	32.0
10.00	45	15.0
12.00	21	7.0
14.00	18	6.0
16.00	18	6.0
18.00	2	.7
Total	300	100.0




Figure (20): Number of chemotherapy cycles among a sample of breast cancer patients

4.4.4 Types of surgery:

Surgical procedures were undertaken for 71.4% of breast cancer patients. Total mastoidectomy was the most common surgery (63.7%) while lobectomy was carried out for 7.7% patients. The distribution of the sample is represented in **table (21)** and graphically illustrated in **Figure (21)**.

Table (21): Types of surgery undertaken for a sample of breast cancer patients:

	Frequency	Percent
no surgery	86	28.7
lopectomy	23	7.7
mastoidectomy	191	63.7
Total	300	100.0




Figure (21): Types of surgery undertaken for a sample of breast cancer patients

4.4.5 Hormonal therapy:

Almost all patients (44%) with estrogen and progesterone receptors positive breast cancer had received any hormonal therapy according to their receptor status. The distribution of the sample is represented in **table (22)** and graphically illustrated in **Figure (22)**.

Table (22): Hormonal therapy among a sample of breast cancer patients:

			Estrogen and progesterone receptor		Total
			negative	positive	
Hormonal therapy	no	Count	168	0	168
		% within Estrogen and progesterone receptor	100.0%	.0%	56.0%
	yes	Count	0	132	132
		% within Estrogen and progesterone receptor	.0%	100.0%	44.0%
Total		Count	168	132	300
		% within Estrogen and progesterone receptor	100.0%	100.0%	100.0%




Figure (22): Hormonal therapy among a sample of breast cancer patients

4.4.6 Biological therapy:

About 42.6% of patients with human epidermal growth factor receptor-2 (HER-2) positive breast cancer have not received biological therapy despite their positive receptor status. The distribution of the sample is represented in **table (23)** and graphically illustrated in **Figure (23)**.

Table (23): Biological therapy among a sample of breast cancer patients:

			HER-2 receptor		Total
			negative	positive	
Trastuzumab therapy	no	Count	145	66	211
		% within HER-2 receptor	100.0%	42.6%	70.3%
	yes	Count	0	89	89
		% within HER-2 receptor	.0%	57.4%	29.7%
Total	Count	145	155	300	
	% within HER-2 receptor	100.0%	100.0%	100.0%	




Figure (23): Biological therapy among a sample of breast cancer patients

4.5 Treatment outcomes of breast cancer:

4.5.1 Post-radiation therapy complications:

The most common post-radiation therapy complications were cutaneous symptoms among 12.3% of the patients while neurological symptoms represented 4% of cases. The distribution of the sample is represented in **table (24)** and graphically illustrated in **Figure (24)**.

Table (24): Post-radiation therapy complications among a sample of breast cancer patients:

	Frequency	Percent
no	248	82.7
cutaneous	37	12.3
neurological	12	4.0
other	1	.3
hematological and neurological	1	.3
cutaneous and neurological	1	.3
Total	300	100.0

Post-radiation complications




Figure (24): Post-radiation therapy complications among a sample of breast cancer patients

4.5.2 Post-surgical complications:

The most common post-surgical complications were neurological symptoms (4%) and lymphedema (2.7%) or both (1.7%). The distribution of the sample is represented in **table (25)** and graphically illustrated in **Figure (25)**.

Table (25): Post-surgical complications among a sample of breast cancer patients:

	Frequency	Percent
no	275	91.7
lymphedema	8	2.7
neurological	12	4.0
lymphedema + neurological	5	1.7
Total	300	100.0




Figure (25): Post-surgical complications among a sample of breast cancer patients

Chapter: 5

Discussion



5. Discussion

The study assessed treatment outcome among breast cancer patients at National Oncology Center, Sana'a. Most of the patients were in the age range of 30-59 years. This might be a result of increased hormonal activity and tissue responsiveness or use of hormonal contraceptives in this age group. The study also showed that many patients were from Sana'a region. This can be explained by the fact that people in Sana'a and nearby regions have easy access to National Oncology Center to seek diagnosis and treatment.

In this study, 300 histologically proven treated breast cancer patients were included. Based on histologic classification, all cases were found to have invasive carcinoma. This is in line with other studies that showed majority of breast cancer cases to be of invasive type (**Breast cancer treatment guideline 2006; Kakarala et al, 2010; Syed et al, 2011; Tovar et al, 2014; Rahal et al, 2015**). This could probably be due to a result of spread to the axillary lymph nodes.

The result of the study showed that majority of the patients came to National Oncology Center when they were on the third stage of the disease. This might be due to low knowledge of the basic symptoms of breast malignancy and absence of a nearby diagnostic center. Majority of cancers



in developing countries are diagnosed at an advanced stage of disease because of lack of screening and early detection services, as well as limited awareness of early signs and symptoms of cancer among the public and health care providers. Stigma associated with diagnosis of cancer also plays a role in late-stage presentation in most parts of Africa (**Tigeneh et al, 2015**). Thus, one can take note of that women who are diagnosed early and put on treatment is one way of improving breast cancer treatment outcomes. However, a different result was reported from a study done in Brazil , where majority of the cases 56 (93%) had an early stage tumor (**Tovar et al, 2014**). This might be is a result of difference in education, economic, social status as well as health seeking behavior between the study population of the studies.

In the present study, chemotherapy and surgical treatment appear to be a pillar component in the National Oncology Center adult oncology protocol. The most common combined regimens used for treatment of breast cancer were AC→T “doxorubicin + cyclophosphamide → taxol” (34.3%) followed by AC “doxorubicin + cyclophosphamide” (16.3%). FAC regimen “5-fluorouracil + doxorubicin + cyclophosphamide” was prescribed for 14%, and FAC→T regiment “5-fluorouracil + doxorubicin + cyclophosphamide → taxol” was prescribed for 13% of the patients. Studies done in Spain



(Martin et al, 2003), Kenya (Wata et al, 2013) and USA (Anampa et al, 2015) found that grater proportional use of FAC regimen for treatment of breast cancer.

Surgery for breast cancer usually involves breast-conserving surgery (BCS) or mastectomy (Cancer Treatment Facts, 2014-2015). Modified radical mastectomy has traditionally been the standard of care for early-stage invasive breast cancers. Therefore, this general mastectomy practice for majority of them in the National Oncology Center may contribute for patient survival. Similar findings were also found in Lebanon that showed mastectomy rates in Arab countries are high amounting to 79.9%–82% in Egypt, 65% in Oman, 70% among Palestinians, 88% in Syria, and 82.4% in Tunisia (Tfayli et al, 2010). This might be because of mastectomies were performed due to the more advanced nature of their breast cancer, more nodal involvement, or even larger tumor size. However, different results were reported in studies from France, where most of the patients had breast-conserving surgery (Rahal et al, 2015). This might be because of histologic subtypes other than invasive ductal carcinoma or acceptable cosmetic outcome can be achieved in almost all patients undergoing breast-conserving surgery without compromising the of local tumor control.



Chapter: 6

Summary and Conclusions



6. Summary and Conclusions

The present study was aimed to assess treatment outcome among breast cancer patients at the National Oncology Cancer. A hospital-based prospective cross-sectional study was conducted by collecting data from medical records of breast cancer patients who attended the breast cancer unit at the National Oncology Center in Sana'a during the period of November 1, 2019 to February 29, 2020.

Total number of sample enrolled in the present study was 300 patients with breast cancer. Patients' age ranged between 21 and 80 years. The present study observed that most cases of breast cancer patients aged between 30-59 years (78.4%). Most patients of breast cancer (31.3%) were from Sana'a and from Taiz (15.7%) governorates. History of contraceptive use was found among 42.3% of breast cancer patients. Age at first menarche was less than 11 years in about 10.3% of breast of breast cancer patients while 89.7% of the patients aged at first menarche between 12 and 18 years. About 18% of breast cancer patients were nulliparous as the either had not children despite marriage or because they are single. Age at first delivery ranged between 12 and 42 years. Most patients 71.3% had got their children while they aged 12-29 years old. The number of children ranged between 1 and 14 for breast cancer women.



Out of all patients, 40.3% % were estrogen positive, 36% were progesterone positive and 51.7% were HER-2 positive. All patients have invasive form of breast cancer. Most of the patients (95%) have ductal invasive breast cancer as compared to 1.7% who have invasive lobular carcinoma. Some patients (2%) have both ductal and lobular invasive breast cancer. About 39% of the breast cancer patients were at stage 3, while 27.7% were at stage 4 and 25.4% were at stage 2.

The most common treatment for breast cancer among the included breast cancer patients was combined surgery and chemotherapy (39%), then the standard method with surgery, radiotherapy, and chemotherapy in 32.7% of the patients. Chemotherapy alone represented 24.7% of the patients while chemotherapy along with radiotherapy represented 3.7% of the patients. The most common combined regimens used for treatment of breast cancer were AC→T “doxorubicin + cyclophosphamide → taxol” (34.3%) followed by AC “doxorubicin + cyclophosphamide” (16.3%). FAC regimen “5-flurouracil + doxorubicin + cyclophosphamide” was prescribed for 14%, and FAC→T regiment “5-flurouracil + doxorubicin + cyclophosphamide → taxol” was prescribed for 13% of the patients. Chemotherapy cycles ranged from 4-18 cycles. Most patients (32%) have received 8 cycles of chemotherapy regimens while other patients received either 6 (17.7%) or 4



cycles (15.7%). Surgical procedures were undertaken for 71.4% of breast cancer patients. Total mastectomy was the most common surgery (63.7%) while lobectomy was carried out for 7.7% patients. Almost all patients with estrogen and progesterone receptors positive breast cancer had received hormonal therapy while 42.6% of patients with human epidermal growth factor receptor-2 (HER-2) positive breast cancer have not received biological therapy despite their positive receptor status.

From the results of the present study, it could be concluded that:

- Most breast cancer patients had estrogen positive (40.3%), progesterone positive (36%), and HER-2 positive (51.7%) disease.
- All patients have invasive form of breast cancer which denotes that patients are usually diagnosed lately.
- Large proportion (42.6%) with human epidermal growth factor receptor-2 (HER-2) positive breast cancer patients have not received biological therapy despite their positive receptor status.

Thus, from these results and conclusions, it could be concluded that:

- National screening of breast cancer is advised especially for patients with risk factors.
- Government is advised to provide the biological therapy for patients with HER-2 positive breast cancer.



Chapter: 7

References




7. References

1. Allred DC, Harvey JM, Berardo M, Clark GM. Prognostic and predictive factors in breast cancer by immunohistochemical analysis. *Mod Pathol* 1998; 11: 155-168.
2. American College of Radiology. Breast imaging reporting and data system (BI-RADS). 4th ed. Reston (VA): American College of Radiology, 2003
3. Anampa J, Makower D, Sparano JA. Progress in adjuvant chemotherapy for breast cancer: an overview. *Spotlight Breast Cancer* 2015; 13:195.
4. Anderson BO, Yip CH, Smith RA, Shyyan R, Sener SF, Eniu A, Carlson RW, Azavedo E, Harford J. Guideline implementation for breast healthcare in low-income and middleincome countries: overview of the Breast Health Global Initiative Global Summit 2007. *Cancer* 2008; 113: 2221-2243.
5. Anderson GL, Manson J, Wallace R, Lund B, Hall D, Davis S, Shumaker S, Wang CY, Stein E, Prentice RL. Implementation of the Women's Health Initiative study design. *Ann Epidemiol* 2003; 13: S5-17.
6. Arafat Tfayli, Sally Temraz, Rachel AbouMrad R. Shamseddine A. Breast Cancer in Low- and Middle-Income Countries: An Emerging and Challenging Epidemic. *J Oncol.* 2010;2010: 490631.
7. Berg WA, Blume JD, Cormack JB, Mendelson EB, Lehrer D, Böhm-Vélez M, Pisano ED, Jong RA, Evans WP, Morton MJ, Mahoney MC, Larsen LH, Barr RG, Farria DM, Marques HS, Boparai K. Combined screening with ultrasound and mammography vs mammography alone in women at elevated risk of breast cancer. *JAMA* 2008; 299: 2151-2163.
8. Berry DA, Cronin KA, Plevritis SK, Fryback DG, Clarke L, Zelen M, Mandelblatt JS, Yakovlev AY, Habbema JD, Feuer EJ. Effect of screening and adjuvant therapy on mortality from breast cancer. *N Engl J Med* 2005; 353: 1784-1792.
9. Breast cancer and hormone replacement therapy: collaborative reanalysis of data from 51 epidemiological studies of 52,705 women with breast cancer and 108,411 women without breast cancer. Collaborative Group on Hormonal Factors in Breast Cancer. *Lancet* 1997; 350: 1047-1059.



10. Buist DS, Abraham LA, Barlow WE, Krishnaraj A, Holdridge RC, Sickles EA, Carney PA, Kerlikowske K, Geller BM. Diagnosis of second breast cancer events after initial diagnosis of early stage breast cancer. *Breast Cancer Res Treat* 2010; 124: 863-873.
11. Cancer Treatment and Survivorship Facts and Figures 2014-2015. Atlanta: American Cancer Society; 2014. Available at : www.cancer.org. Accessed on :01/01/2016.
12. Chen WY, Rosner B, Hankinson SE, Colditz GA, Willett WC. Moderate alcohol consumption during adult life, drinking patterns, and breast cancer risk. *JAMA* 2011; 306: 1884-1890.
13. Chlebowski RT, Manson JE, Anderson GL, Cauley JA, Aragaki AK, Stefanick ML, Lane DS, Johnson KC, Wactawski-Wende J, Chen C, Qi L, Yasmeen S, Newcomb PA, Prentice RL. Estrogen plus progestin and breast cancer incidence and mortality in the Women's Health Initiative Observational Study. *J Natl Cancer Inst* 2013; 105: 526-535.
14. Colditz GA, Kaphingst KA, Hankinson SE, Rosner B. Family history and risk of breast cancer: nurses' health study. *Breast Cancer Res Treat* 2012; 133: 1097-1104.
15. Colditz GA, Rosner B. Cumulative risk of breast cancer to age 70 years according to risk factor status: data from the Nurses' Health Study. *Am J Epidemiol* 2000; 152: 950-964.
16. Coleman MP, Quaresma M, Berrino F, Lutz JM, De Angelis R, Capocaccia R, Baili P, Rachet B, Gatta G, Hakulinen T, Micheli A, Sant M, Weir HK, Elwood JM, Tsukuma H, Koifman S, E Silva GA, Francisci S, Santaquilani M, Verdecchia A, Storm HH, Young JL. Cancer survival in five continents: a worldwide population-based study (CONCORD). *Lancet Oncol* 2008; 9: 730-756.
17. Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and breastfeeding: collaborative reanalysis of individual data from 47 epidemiological studies in 30 countries, including 50302 women with breast cancer and 96973 women without the disease. *Lancet* 2002; 360: 187-195.
18. Collaborative Group on Hormonal Factors in Breast Cancer. Familial breast cancer: collaborative reanalysis of individual data from 52 epidemiological studies including




- 58,209 women with breast cancer and 101,986 women without the disease. *Lancet* 2001; 358: 1389-1399.
19. Curtis RE, Freedman DM, Ron E, Ries LAG, Hacker DG, Edwards BK, Tucker MA, Fraumeni JF Jr., editors. *New Malignancies Among Cancer Survivors: SEER Cancer Registries, 1973-2000*. NIH: National Cancer Institute, 2006
20. Danaei G, Vander Hoorn S, Lopez AD, Murray CJ, Ezzati M. Causes of cancer in the world: comparative risk assessment of nine behavioural and environmental risk factors. *Lancet* 2005; 366: 1784-1793.
21. Di Leo A, Curigliano G, Di-eras V, Malorni L, Sotiriou C, Swanton C, et al (2015). New approaches for improving outcomes in breast cancer in Europe. *The Breast* 24: 332-330.
22. Dipiro JT, Talbert RL, Yee GC, Matzke GR, Wells BG, Posey LM (2011). *Pharmacotherapy: A pathophysiologic approach*. 8edn. McGraw-Hill, New York.
23. Dupont WD, Parl FF, Hartmann WH, Brinton LA, Winfield AC, Worrell JA, Schuyler PA, Plummer WD. Breast cancer risk associated with proliferative breast disease and atypical hyperplasia. *Cancer* 1993; 71: 1258-1265.
24. EMRO Breast cancer treatment guideline. 2006; Available at <www.emro.int/dsaf/dsa697.pdf> (accessed on 01/01/2019).
25. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 2010; 127: 2893-2917.
26. Fitzgibbons PL, Page DL, Weaver D, Thor AD, Allred DC, Clark GM, Ruby SG, O'Malley F, Simpson JF, Connolly JL, Hayes DF, Edge SB, Lichter A, Schnitt SJ. Prognostic factors in breast cancer. College of American Pathologists Consensus Statement 1999. *Arch Pathol Lab Med* 2000; 124: 966-978.
27. Ford D, Easton DF, Stratton M, Narod S, Goldgar D, Devilee P, Bishop DT, Weber B, Lenoir G, Chang-Claude J, Sobol H, Teare MD, Struwing J, Arason A, Scherneck S, Peto J, Rebbeck TR, Tonin P, Neuhausen S, Barkardottir R, Eyfjord J, Lynch H, Ponder BA, Gayther SA, Zelada-Hedman M. Genetic heterogeneity and penetrance analysis of the BRCA1 and BRCA2 genes in breast cancer families. The Breast Cancer Linkage Consortium. *Am J Hum Genet* 1998; 62: 676-689.



28. Freedman DA, Petitti DB, Robins JM. On the efficacy of screening for breast cancer. *Int J Epidemiol* 2004; 33: 43-55.
29. Gartlehner G, Thaler K, Chapman A, Kaminski-Hartenthaler A, Berzaczy D, Van Noord MG, Helbich TH. Mammography in combination with breast ultrasonography versus mammography for breast cancer screening in women at average risk. *Cochrane Database Syst Rev* 2013; 4: CD009632.
30. Gotzsche PC, Jørgensen KJ. Screening for breast cancer with mammography. *Cochrane Database Syst Rev* 2013; 6: CD001877.
31. Guibout C, Adjadj E, Rubino C, Shamsaldin A, Grimaud E, Hawkins M, Mathieu MC, Oberlin O, Zucker JM, Panis X, Lagrange JL, Daly-Schveitzer N, Chavaudra J, de Vathaire F. Malignant breast tumors after radiotherapy for a first cancer during childhood. *J Clin Oncol* 2005; 23: 197-204.
32. Gunter MJ, Hoover DR, Yu H, Wassertheil-Smoller S, Rohan TE, Manson JE, Li J, Ho GY, Xue X, Anderson GL, Kaplan RC, Harris TG, Howard BV, Wylie-Rosett J, Burk RD, Strickler HD. Insulin, insulin-like growth factor-I, and risk of breast cancer in postmenopausal women. *J Natl Cancer Inst* 2009; 101: 48-60.
33. Hammond ME, Hayes DF, Dowsett M, Allred DC, Hagerty KL, Badve S, Fitzgibbons PL, Francis G, Goldstein NS, Hayes M, Hicks DG, Lester S, Love R, Mangu PB, McShane L, Miller K, Osborne CK, Paik S, Perlmutter J, Rhodes A, Sasano H, Schwartz JN, Sweep FC, Taube S, Torlakovic EE, Valenstein P, Viale G, Visscher D, Wheeler T, Williams RB, Wittliff JL, Wolff AC. American Society of Clinical Oncology/ College of American Pathologists guideline recommendations for immunohistochemical testing of estrogen and progesterone receptors in breast cancer (unabridged version). *Arch Pathol Lab Med* 2010; 134: e48-e72.
34. Hartmann LC, Sellers TA, Frost MH, Lingle WL, Degnim AC, Ghosh K, Vierkant RA, Maloney SD, Pankratz VS, Hillman DW, Suman VJ, Johnson J, Blake C, Tlsty T, Vachon CM, Melton LJ, Visscher DW. Benign breast disease and the risk of breast cancer. *N Engl J Med* 2005; 353: 229-237.
35. Henderson TO, Amsterdam A, Bhatia S, Hudson MM, Meadows AT, Neglia JP, Diller LR, Constone LS, Smith RA, Mahoney MC, Morris EA, Montgomery LL, Landier W, Smith SM, Robison LL, Oeffinger KC. Systematic review: surveillance for breast



- cancer in women treated with chest radiation for childhood, adolescent, or young adult cancer. *Ann Intern Med* 2010; 152: 444-455; W144-54.
36. Hicks DG, Kulkarni S. HER2+ breast cancer: review of biologic relevance and optimal use of diagnostic tools. *Am J Clin Pathol* 2008; 129: 263-273.
 37. Hsieh CC, Trichopoulos D, Katsouyanni K, Yuasa S. Age at menarche, age at menopause, height and obesity as risk factors for breast cancer: associations and interactions in an international case-control study. *Int J Cancer* 1990; 46: 796-800.
 38. Kakarala M, Rozek L, Cote M, et al. Breast cancer histology and receptor status characterization in Asian Indian and Pakistani women in the U.S. SEER analysis. *Bio Med Central* 2010; 10:191.
 39. Kelly KM, Dean J, Comulada WS, Lee SJ. Breast cancer detection using automated whole breast ultrasound and mammography in radiographically dense breasts. *Eur Radiol* 2010; 20: 734-742.
 40. Kelsey JL, Gammon MD, John EM. Reproductive factors and breast cancer. *Epidemiol Rev* 1993; 15: 36-47.
 41. Kösters JP, Göttsche PC. Regular self-examination or clinical examination for early detection of breast cancer. *Cochrane Database Syst Rev* 2003; (2): CD003373.
 42. Kriege M, Brekelmans CT, Boetes C, Besnard PE, Zonderland HM, Obdeijn IM, Manoliu RA, Kok T, Peterse H, Tilanus-Linthorst MM, Muller SH, Meijer S, Oosterwijk JC, Beex LV, Tollenaar RA, de Koning HJ, Rutgers EJ, Klijn JG. Efficacy of MRI and mammography for breast-cancer screening in women with a familial or genetic predisposition. *N Engl J Med* 2004; 351: 427-437.
 43. Lahmann PH, Hoffmann K, Allen N, van Gils CH, Khaw KT, Tehard B, Berrino F, Tjønneland A, Bigaard J, Olsen A, Overvad K, Clavel-Chapelon F, Nagel G, Boeing H, Trichopoulos D, Economou G, Bellos G, Palli D, Tumino R, Panico S, Sacerdote C, Krogh V, Peeters PH, Bueno-de-Mesquita HB, Lund E, Ardanaz E, Amiano P, Pera G, Quirós JR, Martínez C, Tormo MJ, Wirfält E, Berglund G, Hallmans G, Key TJ, Reeves G, Bingham S, Norat T, Biessy C, Kaaks R, Riboli E. Body size and breast cancer risk: findings from the European Prospective Investigation into Cancer And Nutrition (EPIC). *Int J Cancer* 2004; 111: 762-771.
 44. Lalloo F, Evans DG. Familial breast cancer. *Clin Genet* 2012; 82: 105-114.



45. Lynch HT, Lynch JF. Breast cancer genetics in an oncology clinic: 328 consecutive patients. *Cancer Genet Cytogenet* 1986; 22: 369-371.
46. Margolin S, Johansson H, Rutqvist LE, Lindblom A, Fornander T. Family history, and impact on clinical presentation and prognosis, in a population-based breast cancer cohort from the Stockholm County. *Fam Cancer* 2006; 5: 309-321.
47. Martin M, Villar A, Sole-Calvo A, et al. Doxorubicin in combination with fluorouracil and cyclophosphamide (i.v. FAC regimen, day 1, 21) versus methotrexate in combination with fluorouracil and cyclophosphamide (i.v. CMF regimen, day 1, 21) as adjuvant chemotherapy for operable breast cancer: a study by the GEICAM group. *Ann Oncol* 2003;14:833–842.
48. McCready T, Littlewood D, Jenkinson J. Breast selfexamination and breast awareness: a literature review. *J Clin Nurs* 2005; 14: 570-578.
49. Milazzo G, Giorgino F, Damante G, Sung C, Stampfer MR, Vigneri R, Goldfine ID, Belfiore A. Insulin receptor expression and function in human breast cancer cell lines. *Cancer Res* 1992; 52: 3924-3930.
50. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology v.1.2013. Breast Cancer Screening and Diagnosis. Available from: URL: [http:// www.nccn.org/professionals/physician_gls/pdf/breast. pdf](http://www.nccn.org/professionals/physician_gls/pdf/breast.pdf) Accessed: December 8, 2013
51. Nyström L, Andersson I, Bjurstam N, Frisell J, Nordenskjöld B, Rutqvist LE. Long-term effects of mammography screening: updated overview of the Swedish randomised trials. *Lancet* 2002; 359: 909-919.
52. Paik S, Shak S, Tang G, Kim C, Baker J, Cronin M, Baehner FL, Walker MG, Watson D, Park T, Hiller W, Fisher ER, Wickerham DL, Bryant J, Wolmark N. A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N Engl J Med* 2004; 351: 2817-2826.
53. Preston DL, Ron E, Tokuoka S, Funamoto S, Nishi N, Soda M, Mabuchi K, Kodama K. Solid cancer incidence in atomic bomb survivors: 1958-1998. *Radiat Res* 2007; 168: 1-64.



54. Pukkala E, Kesminiene A, Poliakov S, Ryzhov A, Drozdovitch V, Kovgan L, Kyyrönen P, Malakhova IV, Gulak L, Cardis E. Breast cancer in Belarus and Ukraine after the Chernobyl accident. *Int J Cancer* 2006; 119: 651-658.
55. Rahal S, Bohe JM, Extra JM, et al. Immunohistochemical subtypes predict the clinical outcome in high-risk node-negative breast cancer patients treated with adjuvant FEC regimen: results of a single-center retrospective study. *Bio Med Central* 2015; 15:697.
56. Ries LAG, Melbert D, Krapcho M, Stinchcomb DG, Howlader N, Horner MJ, Mariotto A, Miller BA, Feuer EJ, Altekruse SF, Lewis DR, Clegg L, Eisner MP, Reichman M, Edwards BK. SEER Cancer Statistics Review, 1975-2005, National Cancer Institute. Available from: URL: http://seer.cancer.gov/csr/1975_2005/
57. Ritte R, Lukanova A, Berrino F, Dossus L, Tjønneland A, Olsen A, Overvad TF, Overvad K, Clavel-Chapelon F, Fournier A, Fagherazzi G, Rohrmann S, Teucher B, Boeing H, Aleksandrova K, Trichopoulou A, Lagiou P, Trichopoulos D, Palli D, Sieri S, Panico S, Tumino R, Vineis P, Quirós JR, Buckland G, Sánchez MJ, Amiano P, Chirlaque MD, Ardanaz E, Sund M, Lenner P, Bueno-de-Mesquita B, van Gils CH, Peeters PH, Krum-Hansen S, Gram IT, Lund E, Khaw KT, Wareham N, Allen NE, Key TJ, Romieu I, Rinaldi S, Siddiq A, Cox D, Riboli E, Kaaks R. Adiposity, hormone replacement therapy use and breast cancer risk by age and hormone receptor status: a large prospective cohort study. *Breast Cancer Res* 2012; 14: R76.
58. Rosner B, Colditz GA, Willett WC. Reproductive risk factors in a prospective study of breast cancer: the Nurses' Health Study. *Am J Epidemiol* 1994; 139: 819-835.
59. Seal S, Thompson D, Renwick A, Elliott A, Kelly P, Barfoot R, Chagtai T, Jayatilake H, Ahmed M, Spanova K, North B, McGuffog L, Evans DG, Eccles D, Easton DF, Stratton MR, Rahman N. Truncating mutations in the Fanconi anemia J gene *BRIP1* are low-penetrance breast cancer susceptibility alleles. *Nat Genet* 2006; 38: 1239-1241.
60. Shapiro S, Strax P, Venet L. Periodic breast cancer screening in reducing mortality from breast cancer. *JAMA* 1971; 215: 1777-1785.
61. Shapiro S, Venet W, Strax P, Venet L, Roeser R. Selection, follow-up, and analysis in the Health Insurance Plan Study: a randomized trial with breast cancer screening. *Natl Cancer Inst Monogr* 1985; 67: 65-74 [PMID: 4047153]



62. Sharif S, Moran A, Huson SM, Iddenden R, Shenton A, Howard E, Evans DG. Women with neurofibromatosis 1 are at a moderately increased risk of developing breast cancer and should be considered for early screening. *J Med Genet* 2007; 44: 481-484.
63. Siegel R, Naishadham D, Jemal A. Cancer statistics, 2013. *CA Cancer J Clin* 2013; 63: 11-30.
64. Sieri S, Krogh V, Bolelli G, Abagnato CA, Grioni S, Pala V, Evangelista A, Allemani C, Micheli A, Tagliabue G, Schunemann HJ, Menard S, Berrino F, Muti P. Sex hormone levels, breast cancer risk, and cancer receptor status in postmenopausal women: the ORDET cohort. *Cancer Epidemiol Biomarkers Prev* 2009; 18: 169-176.
65. Smetherman DH. Screening, imaging, and image-guided biopsy techniques for breast cancer. *Surg Clin North Am* 2013; 93: 309-327.
66. Syed BM, Khyatt WA, Johnston SJ, et al. Long-term clinical outcome of oestrogen receptor-positive operable primary breast cancer in older women: a large series from a single centre. *Br J Cancer* 2011;104:1393-1400.
67. Thompson D, Duedal S, Kirner J, McGuffog L, Last J, Reiman A, Byrd P, Taylor M, Easton DF. Cancer risks and mortality in heterozygous ATM mutation carriers. *J Natl Cancer Inst* 2005; 97: 813-822.
68. Tigeneh W, Molla A, Abreha A, et al. Pattern of cancer in Tikur Anbessa Specialized Hospital Oncology Center in Ethiopia from 1998 to 2010. *Int J Cancer Res Mol Mech* 2015; 1:1-5.
69. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A (2015) Global cancer statistics. *CA Cancer J Clin*. 65(2):87-108.
70. Tovar JR, Zandonade E, Costa Amorim MH. Factors associated with the incidence of local recurrences of breast cancer in women who underwent conservative surgery. *Int J Breast Cancer* 2014; 10:1155.
71. van de Vijver MJ, He YD, van't Veer LJ, Dai H, Hart AA, Voskuil DW, Schreiber GJ, Peterse JL, Roberts C, Marton MJ, Parrish M, Atsma D, Witteveen A, Glas A, Delahaye L, van der Velde T, Bartelink H, Rodenhuis S, Rutgers ET, Friend SH, Bernards R. A gene-expression signature as a predictor of survival in breast cancer. *N Engl J Med* 2002; 347: 1999-2009.



72. Warner E, Messersmith H, Causer P, Eisen A, Shumak R, Plewes D. Systematic review: using magnetic resonance imaging to screen women at high risk for breast cancer. *Ann Intern Med* 2008; 148: 671-679.
73. Wata ED, Osanjo GO, Oluka M, et al. Predictors of breast cancer treatment outcomes in Kenyan women. *Afr J Pharmacol Therap* 2013;2:109–115.
74. Wolff AC, Hammond ME, Schwartz JN, Hagerty KL, Allred DC, Cote RJ, Dowsett M, Fitzgibbons PL, Hanna WM, Langer A, McShane LM, Paik S, Pegram MD, Perez EA, Press MF, Rhodes A, Sturgeon C, Taube SE, Tubbs R, Vance GH, van de Vijver M, Wheeler TM, Hayes DF. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for human epidermal growth factor receptor 2 testing in breast cancer. *J Clin Oncol* 2007; 25: 118-145.
75. Wong MW, Nordfors C, Mossman D, Pecenpetelovska G, Avery-Kiejda KA, Talseth-Palmer B, Bowden NA, Scott RJ. BRIP1, PALB2, and RAD51C mutation analysis reveals their relative importance as genetic susceptibility factors for breast cancer. *Breast Cancer Res Treat* 2011; 127: 853-859.
76. Wu Y, Zhang D, Kang S. Physical activity and risk of breast cancer: a meta-analysis of prospective studies. *Breast Cancer Res Treat* 2013; 137: 869-882.



Chapter: 8

Arabic Abstract



ملخص عربي

معالجة سرطان الثدي في المركز الوطني للأورام – صنعاء : دراسة وصفية

مقدمة: هدفت الدراسة الحالية إلى وصف أنماط العلاج لدى مرضى سرطان الثدي في المركز الوطني للأورام في صنعاء.

الطريقة: أجريت دراسة مستعرضة مستقبلية من خلال جمع البيانات من السجلات الطبية لمرضى سرطان الثدي الذين حضروا وحدة سرطان الثدي في المركز الوطني للأورام في صنعاء خلال الفترة من 1 نوفمبر 2019 إلى 29 فبراير ، 2020.

النتائج: كان إجمالي عدد المرضى المصابين بسرطان الثدي 300 شخصاً معظمهم من محافظتي محافظتي صنعاء وتعز لدى بعضهم (42.3%) تاريخ سابق لاستخدام وسائل منع الحمل. وقد كان بين جميع مرضى سرطان الثدي الذين شملتهم الدراسة 40.3% من الحالات الموجبة لمستقبلات الاستروجين، و 36% موجبة لمستقبلات البروجسترون، بينما 51.7% من الحالات الموجبة لمستقبلات عامل نمو البشرة HER-2. وقد تم تشخيص جميع المرضى بأن لديهم شكل من أشكال سرطان الثدي الغازية. وكان معظم المرضى (95%) يعانون من سرطان الثدي الغازي القنوات. وتم تشخيص 39 % من مرضى سرطان الثدي في المرحلة الثالثة ، بينما كان 27.7 % في المرحلة الرابعة و 25.4 % في المرحلة الثانية. وكان العلاج الأكثر شيوعاً لسرطان الثدي بين المرضى هي الجراحة مع العلاج الكيميائي (39 %) ثم طريقة العلاج بالجراحة مع العلاج الإشعاعي والعلاج الكيميائي في 32.7% من المرضى. وكان العلاج الكيماوي وحده يمثل 24.7% من المرضى بينما العلاج الكيماوي مع العلاج الإشعاعي يمثل 3.7% من المرضى. وتم إجراء العمليات الجراحية لـ



71.4% من مرضى سرطان الثدي. وظهرت النتائج أن 42.6% من المرضى الذين يعانون من سرطان الثدي الإيجابي لمستقبلات عامل نمو البشرة HER-2 لم يتلقوا علاجًا بيولوجيًا على الرغم من حالة مستقبلاتهم الإيجابية.

الاستنتاجات: جميع المرضى يعانون من شكل سرطان الثدي الغازية مما يدل على أن المرضى عادة ما يتم تشخيصهم مؤخرًا. وبالتالي ، ينصح بالكشف عن سرطان الثدي خاصة للمرضى الذين يعانون من عوامل الخطر. ونسبة كبيرة (42.6%) من المرضى الذين لديهم سرطان الثدي إيجابي لمستقبلات عامل نمو البشرة HER-2 لم يتلقوا العلاج البيولوجي على الرغم من حالة مستقبلاتهم الإيجابية.





الجمهورية اليمنية
وزارة التعليم العالي والبحث العلمي
الجامعة الاماراتية الدولية
كلية الطب والعلوم الصحية
قسم الصيدلة السريرية

معالجة سرطان الثدي في المركز الوطني للأورام – صنعاء: دراسة وصفية

بحث تخرج مقدم كمتمم جزئي للحصول على درجة البكالوريوس في الصيدلة السريرية

مقدم من الطلاب:

عبدالكافي عبدالله عبدالكريم محمد

محمد منير حميد

نايف محمد العامري

زايد احمد بعموم

عبدالحاميد احمد محمد الجندبي

امل علي هادي عيسى

فاطمة خالد محمد ابوبكر

احمد علي احمد منصور الخبائي

شعيب كامل درهم ناجي

تحت اشراف:

د/ شوقي حسين ناجي العودي

أستاذ مساعد علم الأدوية والصيدلة السريرية

كلية الطب والعلوم الصحية – جامعة ذمار

د/ عبد الله ثوابه

أستشاري أورام وعلاج اشعاعي

مدير المركز الوطني للأورام – صنعاء

2020-2019

